



THE UNIVERSITY *of* EDINBURGH

Edinburgh Research Explorer

A derived equivalence for a degree 6 del Pezzo surface over an arbitrary field

Citation for published version:

Blunk, M, Sierra, SJ & Smith, SP 2011, 'A derived equivalence for a degree 6 del Pezzo surface over an arbitrary field', *Journal of k-Theory*, vol. 8, no. 3, pp. 481-492. <https://doi.org/10.1017/is010011013jkt134>

Digital Object Identifier (DOI):

[10.1017/is010011013jkt134](https://doi.org/10.1017/is010011013jkt134)

Link:

[Link to publication record in Edinburgh Research Explorer](#)

Document Version:

Early version, also known as pre-print

Published In:

Journal of k-Theory

Publisher Rights Statement:

Copyright © ISOPP 2010

DOI: <http://dx.doi.org/10.1017/is010011013jkt134>

General rights

Copyright for the publications made accessible via the Edinburgh Research Explorer is retained by the author(s) and / or other copyright owners and it is a condition of accessing these publications that users recognise and abide by the legal requirements associated with these rights.

Take down policy

The University of Edinburgh has made every reasonable effort to ensure that Edinburgh Research Explorer content complies with UK legislation. If you believe that the public display of this file breaches copyright please contact openaccess@ed.ac.uk providing details, and we will remove access to the work immediately and investigate your claim.



A DERIVED EQUIVALENCE FOR A DEGREE 6 DEL PEZZO SURFACE OVER AN ARBITRARY FIELD

M. BLUNK, S.J. SIERRA, AND S. PAUL SMITH

ABSTRACT. Let S be a degree six del Pezzo surface over an arbitrary field F . Motivated by the first author's classification of all such S up to isomorphism [3] in terms of a separable F -algebra $B \times Q \times F$, and by his K-theory isomorphism $K_n(S) \cong K_n(B \times Q \times F)$ for $n \geq 0$, we prove an equivalence of derived categories

$$D^b(\text{coh}S) \equiv D^b(\text{mod}A)$$

where A is an explicitly given finite dimensional F -algebra whose semisimple part is $B \times Q \times F$.

1. INTRODUCTION

We will work over an arbitrary field F .

Throughout S denotes a degree six del Pezzo surface over F . Equivalently, S is a smooth projective surface over F whose anti-canonical sheaf is ample and has self-intersection number 6.

Throughout \bar{F} will denote a separable closure of F and we will write

$$\bar{S} = S_{\bar{F}} = S \times_{\text{Spec } F} \text{Spec } \bar{F}.$$

In [3], the first author classified such S up to isomorphism by associating to S a pair of separable F -algebras B and Q , both defined as endomorphism rings of certain locally free sheaves on S . Furthermore, it was shown there that the algebraic K-theory of S is isomorphic to that of the algebra $B \times Q \times F$.

Let $\text{coh}S$ denote the category of coherent sheaves on S and let $\text{mod}A$ denote the category of noetherian right A -modules. Let \equiv denote equivalence of derived categories. Our main result (Theorem 4.5) establishes a derived equivalence

$$(1-1) \quad D^b(\text{coh}S) \equiv D^b(\text{mod}A)$$

where A is a finite dimensional F -algebra whose semi-simple quotient is $B \times Q \times F$. We prove this equivalence by constructing a tilting bundle \mathcal{T} on S that has A as its endomorphism ring. (The definition of a tilting bundle is given in section 4.) The main novelty of our approach is that we do not make any assumptions on the base field F . Since the field F is arbitrary, we cannot assume that S is obtained by blowing up \mathbb{P}_F^2 (in fact S could be a minimal surface), nor can we use exceptional collections.

1991 *Mathematics Subject Classification.* 11E35, 14F05, 14J26, 14J60.

Key words and phrases. Derived equivalence, del Pezzo surface, arbitrary field.

M. Blunk was supported by the National Science Foundation, Award No. 0902659. S.J. Sierra was supported by the National Science Foundation, Award No. 0802935. S.P. Smith was supported by the National Science Foundation, Award No. 0602347.


Acknowledgments. All three authors acknowledge the support of the National Science Foundation with gratitude. We would also like to thank the referee for their comments. This research was partially done while the first author was visiting the University of Washington, and he would like to thank the institution for its support and excellent working conditions. Finally, the first author would like to thank Aravind Asok, Baptise Calmès, and Daniel Krashen for suggesting this problem.

2. BASIC FACTS ABOUT \bar{S}

In this section, we give basic facts about the degree 6 del Pezzo surface \bar{S} . Since all the results here are well-known, we do not give references.

There are six (-1) -curves on \bar{S} , which we may take to lie in the following configuration:

(2-1)



The Picard group is

$$\text{Pic } \bar{S} \cong \frac{\bigoplus_{i=1}^3 (\mathbb{Z}L_i \oplus \mathbb{Z}M_i)}{(M_i + L_j = M_j + L_i \mid 1 \leq i, j \leq 3)}.$$

Usually we only care about the class of a divisor in $\text{Pic } \bar{S}$. We will write

$$D_1 \sim D_2$$

if D_1 and D_2 are linearly equivalent divisors.

As remarked in the discussion after Prop. 2.1 in [3], the group of connected components of the group $\text{Aut } \bar{S}$ is $S_2 \times S_3$, which can be identified with the automorphism group of the hexagon of (-1) -curves on \bar{S} . In particular, there is an element $\sigma \in \text{Aut}(\bar{S})$ that cyclically permutes the six exceptional lines. It is easy to see that $(1 + \sigma)(1 - \sigma^3)$ acts trivially on $\text{Pic } \bar{S}$.

An anti-canonical divisor is

$$-K_{\bar{S}} := L_1 + L_2 + L_3 + M_1 + M_2 + M_3.$$

This is ample. We define two particular divisors

$$(2-2) \quad H := L_1 + M_2 + M_3 \sim L_2 + M_1 + M_3 \sim L_3 + M_1 + M_2$$

and

$$(2-3) \quad H' := L_1 + L_2 + M_3 \sim L_2 + L_3 + M_1 \sim L_3 + L_1 + M_2$$

on \bar{S} . Note that $\sigma(H) \sim H'$ and $\sigma^2(H) \sim H$.

We define the **degree** of a divisor C on \bar{S} as $\deg C = -C \cdot K$. Each exceptional line has degree 1.

There are two morphisms $f, f' : \bar{S} \rightarrow \mathbb{P}_{\bar{F}}^2$, each of which realizes \bar{S} as the blowup of $\mathbb{P}_{\bar{F}}^2$ at three non-collinear points. We choose these so that f contracts the lines L_1, L_2 , and L_3 and f' contracts the lines M_1, M_2 , and M_3 . These two morphisms induce injective group homomorphisms $f^*, f'^* : \text{Pic } \mathbb{P}^2 \rightarrow \text{Pic } \bar{S}$. If ℓ is a line on $\mathbb{P}_{\bar{F}}^2$, then $f^*\ell = H$ and $f'^*\ell = H'$.

The action of $\text{Gal}(\bar{F}/F)$ on the exceptional lines on \bar{S} induces actions of $\text{Gal}(\bar{F}/F)$ on

$$\bar{\mathcal{I}} := \bigoplus_{i=0}^5 \mathcal{O}_{\bar{S}}(\sigma^i H)$$

and

$$\bar{\mathcal{J}} := \bigoplus_{i=0}^5 \mathcal{O}_{\bar{S}}(\sigma^i(L_1 + M_2))$$

that are compatible with its action on \bar{S} . In particular, $\bar{\mathcal{I}}$ and $\bar{\mathcal{J}}$ are $\text{Gal}(\bar{F}/F)$ -invariant. It follows that the locally free sheaves $\bar{\mathcal{I}}$ and $\bar{\mathcal{J}}$ descend to locally free sheaves \mathcal{I} and \mathcal{J} on S .

Define

$$\bar{\mathcal{T}} := \bar{\mathcal{I}} \oplus \bar{\mathcal{J}} \oplus \mathcal{O}_{\bar{S}}, \quad \mathcal{T} := \mathcal{I} \oplus \mathcal{J} \oplus \mathcal{O}_S,$$

and

$$B := \text{End}_S \mathcal{I}, \quad Q := \text{End}_S \mathcal{J}, \quad A := \text{End}_S \mathcal{T}.$$

In [3] it is shown that S is determined up to isomorphism by the pair of F -algebras (B, Q) . (Actually, in [3], B is defined as $(\text{End}_S \mathcal{I}^\vee)^{\text{op}}$. Since sending a homomorphism $\alpha : \mathcal{I} \rightarrow \mathcal{I}$ to its transpose $\alpha^\vee : \mathcal{I}^\vee \rightarrow \mathcal{I}^\vee$ is an anti-isomorphism from $\text{End}_S \mathcal{I}$ to $\text{End}_S \mathcal{I}^\vee$, our B is the same as that in [3], and similarly for Q .) As discussed in [3], the algebras B and Q are Azumaya over their centers, which are respectively étale quadratic and cubic extensions of F . Moreover, these étale centers can be recovered from the action of $\text{Gal}(\bar{F}/F)$ on the hexagon of (-1) -curves, as the action induces a 1-cocycle of $\text{Gal}(\bar{F}/F)$ with values in $S_2 \times S_3$, inducing a pair of étale extensions of F , quadratic and cubic.

We end this section with two results about the endomorphism algebra of \mathcal{T} .

Lemma 2.1. *Let $A := \text{End}_S \mathcal{T}$. Then*

$$A = \begin{pmatrix} B & \text{Hom}_S(\mathcal{J}, \mathcal{I}) & \text{Hom}_S(\mathcal{O}_S, \mathcal{I}) \\ 0 & Q & \text{Hom}_S(\mathcal{O}_S, \mathcal{J}) \\ 0 & 0 & F \end{pmatrix}.$$

Proof. It suffices to show $\text{Hom}_{\bar{S}}(\bar{\mathcal{I}}, \bar{\mathcal{J}}) = \text{Hom}_{\bar{S}}(\bar{\mathcal{I}}, \mathcal{O}_{\bar{S}}) = \text{Hom}_{\bar{S}}(\bar{\mathcal{J}}, \mathcal{O}_{\bar{S}}) = 0$. However, each of these three Hom-spaces is isomorphic to a direct sum of terms of the form $H^0(\bar{S}, \mathcal{O}_{\bar{S}}(D))$ for a divisor D with $\deg D < 0$. But if D has a section then $D \sim D'$ for some effective D' so $\deg D = -D' \cdot K \geq 0$. These Hom spaces are therefore zero. \square

The projective dimension of a left T -module is denoted by $\text{pdim}_T M$. The global homological dimension of T is defined and denoted by

$$\text{gldim } T := \sup\{\text{pdim}_T M \mid M \in \text{Mod } T\}.$$

Proposition 2.2. $\text{gldim } A \leq 2$.

Proof. Let R and S be rings and X an R - S -bimodule. If S is a semisimple ring, then

$$\text{gldim} \begin{pmatrix} R & X \\ 0 & S \end{pmatrix} = \max\{\text{pdim}_R X + 1, \text{gldim } R\}.$$

(See [1, Prop. III.2.7].) Applying this result twice, first to

$$(2-4) \quad A' := \begin{pmatrix} B & \text{Hom}(\mathcal{J}, \mathcal{I}) \\ 0 & Q \end{pmatrix}$$

then to A with $R = A'$ and $S = F$, gives the desired result. \square

3. COHOMOLOGY VANISHING LEMMAS

We will prove several results about vanishing of cohomology and Ext-groups for sheaves on S . These results will be used in Section 4 to show that \mathcal{T} is a tilting bundle and therefore induces an equivalence of derived categories.

A key step in proving that \mathcal{T} is tilting is showing that $\text{Ext}_S^i(\mathcal{T}, \mathcal{T}) = 0$ for $i > 0$. This reduces, by flat base change, to proving that $\text{Ext}_S^i(\overline{\mathcal{T}}, \overline{\mathcal{T}}) = 0$. Given the explicit description of $\overline{\mathcal{T}}$ as a direct sum of invertible sheaves, it suffices to prove that $h^1(D - D') = h^2(D - D') = 0$ for all D and D' belonging to the list

$$(3-1) \quad H, \quad H', \quad L_1 + M_2, \quad L_2 + M_3, \quad L_3 + M_1, \quad 0.$$

We will make repeated use of the relation $L_i + M_j \sim L_j + M_i$.

Proposition 3.1. *Let D and D' be divisors on \bar{S} appearing in the list (3-1). Then*

$$-3 \leq \deg(D - D') \leq 3.$$

Furthermore,

- (1) *if $\deg(D - D') = 1$, then $D - D'$ is linearly equivalent to an exceptional line.*
- (2) *if $\deg(D - D') = 2$, then $D - D' \sim L_i + M_j$ for some $i \neq j \in \{1, 2, 3\}$.*
- (3) *if $\deg(D - D') = 3$, then $D - D'$ is linearly equivalent to either H or H' .*
- (4) *if $\deg(D - D') = 0$, then $D - D'$ is linearly equivalent to either 0 , $L_i - L_j$, $L_i - M_i$, or $M_i - L_i$ for some $i, j \in \{1, 2, 3\}$.*
- (5) *if $\deg(D - D') < 0$, then $D - D'$ is linearly equivalent to either $-L_i$, or $-M_j$, or $-L_i - M_j$, or $-H$, or $-H'$.*

Proof. Exceptional lines have degree 1 so $\deg H = \deg H' = 3$ and $\deg(L_i + M_j) = 2$. It follows that the degree of $D - D'$ is between 3 and -3.

(1) If $\deg(D - D') = 1$, then D is linearly equivalent to H or H' and $D' = L_i + M_j$ for some i, j . It follows from (2-2) and (2-3) that $D - D'$ is linearly equivalent to an exceptional line, and every exceptional line can occur as $D - D'$.

(2) and (3) are obvious.

(4) In this case D and D' have the same degree.

If $\deg D = \deg D' = 2$, then $D = L_i + M_j$ and $D' = L_k + M_\ell$. By considering all possible i, j, k, ℓ , we see that $D - D'$ is linearly equivalent to a divisor of the form $L_i - L_j$.

If $\deg D = \deg D' = 3$, then, for example, $D \sim H$ and $D' \sim H'$, and $D - D' \sim L_i - M_i$. Switching the roles of H and H' , we see $D - D' \sim M_i - L_i$. Finally, we may have $D - D' \sim 0$.

(5) This is the mirror of the cases (1)-(3). \square

Corollary 3.2. *Suppose D is the difference of two divisors appearing in the list (3-1). If $\deg D \geq -2$, then there is an exceptional line E on \bar{S} such that $D - E$ is also a difference of two divisors appearing in the list (3-1) and $D.E \geq -1$.*

Proof. This is established through case-by-case analysis using Proposition 3.1 to look at all the possibilities for D . \square

A divisor D on \bar{S} is good if $h^1(D) = h^2(D) = 0$.

Lemma 3.3. *The divisors $-H$ and $-H'$ on \bar{S} are good.*

Proof. The existence of the morphisms $f, f' : \bar{S} \rightarrow \mathbb{P}_F^2$ allows us to use the Leray spectral sequence. The arguments for $-H$ and $-H'$ are the same so we only prove the result for $-H$.

Because \bar{S} is a blowup of \mathbb{P}_F^2 , $f_*\mathcal{O}_{\bar{S}} = \mathcal{O}_{\mathbb{P}_F^2}$ and $R^j f_*\mathcal{O}_{\bar{S}} = 0$ if $j \geq 1$.

Since $\mathcal{O}_{\bar{S}}(-H) \cong f^*\mathcal{O}_{\mathbb{P}_F^2}(-\ell)$, the projection formula gives

$$\begin{aligned} R^j f_*\mathcal{O}_{\bar{S}}(-H) &= R^j f_*(\mathcal{O}_{\bar{S}} \otimes f^*\mathcal{O}_{\mathbb{P}_F^2}(-\ell)) \\ &\cong R^j f_*\mathcal{O}_{\bar{S}} \otimes \mathcal{O}_{\mathbb{P}_F^2}(-\ell) \\ &\cong \begin{cases} \mathcal{O}_{\mathbb{P}_F^2}(-\ell) & \text{if } j = 0 \\ 0 & \text{if } j \neq 0. \end{cases} \end{aligned}$$

The Leray spectral sequence

$$H^i(\mathbb{P}_F^2, R^j f_*\mathcal{O}_{\bar{S}}(-H)) \Rightarrow H^{i+j}(\bar{S}, \mathcal{O}_{\bar{S}}(-H))$$

therefore degenerates to give

$$H^i(\bar{S}, \mathcal{O}_{\bar{S}}(-H)) \cong H^i(\mathbb{P}_F^2, \mathcal{O}_{\mathbb{P}_F^2}(-\ell))$$

for all i . The result follows because $H^i(\mathbb{P}_F^2, \mathcal{O}_{\mathbb{P}_F^2}(-\ell)) = 0$ for all i . \square

Lemma 3.4. *Let C be any divisor on \bar{S} , and let E be one of the (-1) -curves. If $C - E$ is good and $C.E \geq -1$, then C is good.*

Proof. The long exact sequence in cohomology associated to

$$0 \rightarrow \mathcal{O}_{\bar{S}}(C - E) \rightarrow \mathcal{O}_{\bar{S}}(C) \rightarrow \mathcal{O}_E(C) \rightarrow 0$$

reads in part

$$\begin{aligned} \longrightarrow H^1(\bar{S}, \mathcal{O}_{\bar{S}}(C - E)) &\longrightarrow H^1(\bar{S}, \mathcal{O}_{\bar{S}}(C)) \longrightarrow H^1(\bar{S}, \mathcal{O}_E(C)) \longrightarrow \\ &\longrightarrow H^2(\bar{S}, \mathcal{O}_{\bar{S}}(C - E)) \longrightarrow H^2(\bar{S}, \mathcal{O}_{\bar{S}}(C)) \longrightarrow H^2(\bar{S}, \mathcal{O}_E(C)). \end{aligned}$$

By hypothesis, the left-most term in each row is zero. The right-most term in each row is also zero because $H^i(\bar{S}, \mathcal{O}_E(C)) \cong H^i(\mathbb{P}_F^1, \mathcal{O}_{\mathbb{P}_F^1}(C.E))$. Hence C is good. \square

4. THE TILTING BUNDLE \mathcal{T}

In this section, we show that \mathcal{T} is a tilting bundle and prove our main result.

Proposition 4.1. *Let $i \geq 1$. Then $\text{Ext}_S^i(\mathcal{T}, \mathcal{T}) = 0$.*

Proof. By flat base change it suffices to prove this when F is separably closed so we assume that $F = \bar{F}$. In that case $\text{Ext}_S^i(\mathcal{T}, \mathcal{T})$ is isomorphic to a direct sum of terms of the form $H^i(S, \mathcal{O}_S(D - D'))$ where D and D' are divisors in the list (3-1).

It therefore suffices to show that $D - D'$ is good whenever D and D' are divisors in the list (3-1).

We argue by induction on $\deg(D - D')$. By Proposition 3.1, $-3 \leq \deg(D - D') \leq 3$. If $\deg(D - D') = -3$, then $D - D'$ is good by Lemma 3.3. Now suppose that $-2 \leq \deg(D - D') \leq 3$. By Corollary 3.2, there is an exceptional line E such that $D - D' - E$ is a difference of divisors in (3-1) and $(D - D') \cdot E \geq -1$. By the induction hypothesis, $D - D' - E$ is good, and it then follows from Lemma 3.4 that $D - D'$ is good. \square

Since \bar{S} is a del Pezzo surface of degree ≥ 6 it is a toric variety so we can, and will, make use of Cox's homogeneous coordinate ring for it [5].

Lemma 4.2. *Every $\mathcal{F} \in \text{coh}\bar{S}$ has a finite resolution in which all terms are direct sums of invertible sheaves $\mathcal{O}_{\bar{S}}(D)$ for various divisors D on \bar{S} .*

Proof. Let A be Cox's homogeneous coordinate ring for \bar{S} [5]. Then A is a polynomial ring with a grading by $\text{Pic}(\bar{S})$. Let M be a finitely generated graded A -module. Then M has a finite projective resolution in the category of graded A -modules. By [9, Lemma 2.2], every finitely generated projective graded A -module is a direct sum of twists of A . The exact functor $\text{Gr}(A, \text{Pic}(\bar{S})) \rightarrow \text{Qcoh}\bar{S}$, $M \mapsto \widetilde{M}$, described in [5, Thm. 3.11] sends the resolution of M to an exact sequence in $\text{Qcoh}\bar{S}$ in which the right-most term is \widetilde{M} and all other terms are direct sums of various $\mathcal{O}_{\bar{S}}(D)$, $D \in \text{Div}(\bar{S})$. Given $\mathcal{F} \in \text{coh}\bar{S}$, there is a finitely generated graded A -module M such that $\mathcal{F} \cong \widetilde{M}$. \square

For the rest of this paper, we will work in the derived category. If \mathcal{D} is a triangulated category, we denote the shift of an object \mathcal{M} by $\mathcal{M}[1]$. Recall that a subcategory of \mathcal{D} is *thick* (épaisse) if it is closed under isomorphisms, shifts, taking cones of morphisms, and taking direct summands of objects.

Let \mathcal{D} be a triangulated category and \mathcal{E} a set of objects in \mathcal{D} . Then

- \mathcal{D}^c denotes the full subcategory of \mathcal{D} consisting of the compact objects, i.e., those objects C such that $\text{Hom}_{\mathcal{D}}(C, -)$ commutes with direct sums;
- $\langle \mathcal{E} \rangle$ denotes the smallest thick full triangulated subcategory of \mathcal{D} containing \mathcal{E} ;
- \mathcal{E}^\perp denotes the full subcategory of \mathcal{D} consisting of objects \mathcal{M} such that $\text{Hom}_{\mathcal{D}}(E[i], \mathcal{M}) = 0$ for all $E \in \mathcal{E}$ and all $i \in \mathbb{Z}$.

We say that

- \mathcal{E} generates \mathcal{D} if $\mathcal{E}^\perp = 0$ and that
- \mathcal{D} is compactly generated if $(\mathcal{D}^c)^\perp = 0$.

Clearly, if \mathcal{D} is compactly generated and $\langle \mathcal{E} \rangle = \mathcal{D}^c$, then \mathcal{E} generates \mathcal{D} .

Theorem 4.3 (Ravenel and Neeman [8]. Also see Thm. 2.1.2 in [4]). *Let \mathcal{D} be a compactly generated triangulated category. Then a set of objects $\mathcal{E} \subset \mathcal{D}^c$ generates \mathcal{D} if and only if $\langle \mathcal{E} \rangle = \mathcal{D}^c$.* \square

The unbounded derived categories $D(\text{Qcoh}S)$ and $D(\text{Qcoh}\bar{S})$ are compactly generated. Moreover, $D(\text{Qcoh}S)^c = D^b(\text{coh}S)$ and $D(\text{Qcoh}\bar{S})^c = D^b(\text{coh}\bar{S})$.

Tilting bundles. Let X be a projective scheme over a field k . A locally free sheaf $\mathcal{T} \in \text{coh}X$ is a tilting bundle if it generates $D(\text{Qcoh}X)$ and $\text{Ext}_X^i(\mathcal{T}, \mathcal{T}) = 0$ for all $i > 0$.

Theorem 4.4. $\bar{\mathcal{T}}$ generates $D(\text{Qcoh}\bar{S})$ and $\langle \bar{\mathcal{T}} \rangle = D^b(\text{coh}\bar{S})$.

Proof. By Theorem 4.3, it suffices to show that $\langle \bar{\mathcal{T}} \rangle = D^b(\text{coh}\bar{S})$. Since $\langle \text{coh}\bar{S} \rangle = D^b(\text{coh}\bar{S})$ it suffices to show that every coherent $\mathcal{O}_{\bar{S}}$ -module belongs to $\langle \bar{\mathcal{T}} \rangle$.

If D is an effective divisor on \bar{S} we write \mathcal{I}_D for the ideal vanishing on D as a scheme. Thus $\mathcal{I}_D \cong \mathcal{O}_{\bar{S}}(-D)$. Whenever we write an arrow $\mathcal{O}_{\bar{S}}(-D) \rightarrow \mathcal{O}_{\bar{S}}$ it will be with the tacit understanding that this is the composition of an isomorphism $\mathcal{O}_{\bar{S}}(-D) \rightarrow \mathcal{I}_D$ followed by the inclusion $\mathcal{I}_D \rightarrow \mathcal{O}_{\bar{S}}$.

Since $M_3 \cdot (L_1 + M_2 + M_3) = 0$, $\mathcal{O}_{M_3} \cong \mathcal{O}_{M_3}(L_1 + M_2 + M_3)$. It follows from the exact sequences

$$0 \rightarrow \mathcal{O}_{\bar{S}}(L_1 + M_2) \rightarrow \mathcal{O}_{\bar{S}}(L_1 + M_2 + M_3) \rightarrow \mathcal{O}_{M_3}(L_1 + M_2 + M_3) \rightarrow 0$$

and

$$0 \rightarrow \mathcal{O}_{\bar{S}}(-M_3) \rightarrow \mathcal{O}_{\bar{S}} \rightarrow \mathcal{O}_{M_3} \rightarrow 0$$

that \mathcal{O}_{M_3} and $\mathcal{O}_{\bar{S}}(-M_3)$ belong to $\langle \bar{\mathcal{T}} \rangle$. Hence \mathcal{O}_E and $\mathcal{O}_{\bar{S}}(-E)$ belong to $\langle \bar{\mathcal{T}} \rangle$ for all exceptional lines E .

Since $L_i \cdot L_k = 0$ if $i \neq k$, there is an exact sequence

$$0 \rightarrow \mathcal{O}_{\bar{S}}(-L_i - L_k) \rightarrow \mathcal{O}_{\bar{S}}(-L_i) \oplus \mathcal{O}_{\bar{S}}(-L_k) \rightarrow \mathcal{O}_{\bar{S}} \rightarrow 0.$$

Twisting by $L_i + M_j + L_k$, we obtain

$$0 \rightarrow \mathcal{O}_{\bar{S}}(M_j) \rightarrow \mathcal{O}_{\bar{S}}(M_j + L_k) \oplus \mathcal{O}_{\bar{S}}(L_i + M_j) \rightarrow \mathcal{O}_{\bar{S}}(L_i + M_j + L_k) \rightarrow 0.$$

Therefore, $\mathcal{O}_{\bar{S}}(M_j) \in \langle \bar{\mathcal{T}} \rangle$. From the exact sequence

$$0 \rightarrow \mathcal{O}_{\bar{S}} \rightarrow \mathcal{O}_{\bar{S}}(M_j) \rightarrow \mathcal{O}_{M_j}(M_j) \rightarrow 0,$$

we deduce that $\mathcal{O}_{M_j}(M_j) \in \langle \bar{\mathcal{T}} \rangle$.

It follows that $\mathcal{O}_E(E) \in \langle \bar{\mathcal{T}} \rangle$ for every exceptional curve E . But \mathcal{O}_E is also in $\langle \bar{\mathcal{T}} \rangle$ so, because $D^b(\text{coh}\mathbb{P}_{\bar{F}}^1)$ is generated by $\mathcal{O}_{\mathbb{P}_{\bar{F}}^1}$ and $\mathcal{O}_{\mathbb{P}_{\bar{F}}^1}(-1)$, it follows that $D^b(\text{coh}E) \subset \langle \bar{\mathcal{T}} \rangle$. Hence $\mathcal{O}_E(D) \in \langle \bar{\mathcal{T}} \rangle$ for all divisors D on \bar{S} .

Suppose $\mathcal{O}_{\bar{S}}(D) \in \langle \bar{\mathcal{T}} \rangle$. Then $\mathcal{O}_{\bar{S}}(D - E) \in \langle \bar{\mathcal{T}} \rangle$ because there is an exact sequence

$$0 \rightarrow \mathcal{O}_{\bar{S}}(D - E) \rightarrow \mathcal{O}_{\bar{S}}(D) \rightarrow \mathcal{O}_E(D) \rightarrow 0.$$

Likewise, $\mathcal{O}_{\bar{S}}(D + E) \in \langle \bar{\mathcal{T}} \rangle$ because there is an exact sequence

$$0 \rightarrow \mathcal{O}_{\bar{S}}(D) \rightarrow \mathcal{O}_{\bar{S}}(D + E) \rightarrow \mathcal{O}_E(D + E) \rightarrow 0.$$

It follows that $\langle \bar{\mathcal{T}} \rangle$ contains $\mathcal{O}_{\bar{S}}(D)$ for all $D \in \text{Div } \bar{S}$ and therefore, by Lemma 4.2, contains \mathcal{F} for every $\mathcal{F} \in \text{coh}\bar{S}$. \square

When F is not separably closed \mathcal{T} need not split as a direct sum of line bundles so the arguments in Theorem 4.4 can not be used to prove directly that $\langle \mathcal{T} \rangle = D^b(\text{coh}S)$. Instead we will show that \mathcal{T} generates $D(\text{Qcoh}S)$ and then apply Theorem 4.3.

Theorem 4.5. *Let F be an arbitrary field. Then*

$$\mathrm{RHom}_S(\mathcal{T}, -) : \mathrm{D}^b(\mathrm{coh}S) \rightarrow \mathrm{D}^b(\mathrm{mod}A)$$

is an equivalence of categories.

Proof. We will show that \mathcal{T} generates $\mathrm{D}(\mathrm{Qcoh}S)$. It will then follow from Theorem 4.3 that

$$\langle \mathcal{T} \rangle = \mathrm{D}(\mathrm{Qcoh}S)^c = \mathrm{D}^b(\mathrm{coh}S).$$

By Proposition 4.1, $\mathrm{Ext}_S^i(\mathcal{T}, \mathcal{T}) = 0$ for $i > 0$. By Proposition 2.2, $A = \mathrm{End}_S(\mathcal{T})$ has finite global dimension. Thus we have shown that \mathcal{T} is a tilting bundle and our theorem will then follow directly from [2, Thm. 3.1.2] (or [6, Thm. 7.6]).

Let $\mathcal{M} \in \mathrm{D}(\mathrm{Qcoh}S)$ and suppose $\mathrm{RHom}_S(\mathcal{T}, \mathcal{M}) = 0$. We must show that $\mathcal{M} = 0$.

Since \mathcal{T} is locally free, $\mathcal{H}om_S(\mathcal{T}, -)$ and $\mathcal{T}^\vee \otimes_S -$ are exact functors on $\mathrm{Qcoh}S$. Likewise, $\mathcal{H}om_{\bar{S}}(\bar{\mathcal{T}}, -)$ and $\bar{\mathcal{T}}^\vee \otimes_{\bar{S}} -$ are exact functors on $\mathrm{Qcoh}\bar{S}$. Thus, for example, $\mathrm{RHom}_S(\mathcal{T}, \mathcal{M})$ can be computed on $\mathrm{D}(\mathrm{Qcoh}S)$ by applying $\mathcal{H}om_S(\mathcal{T}, -)$ to each individual term in \mathcal{M} .

Consider the cartesian square

$$\begin{array}{ccc} \bar{S} & \xrightarrow{v} & S \\ q \downarrow & & \downarrow p \\ \mathrm{Spec}(\bar{F}) & \xrightarrow{u} & \mathrm{Spec}(F). \end{array}$$

Since u (and therefore v) is flat, the natural transformation

$$u^* R p_* \rightarrow R q_* v^*$$

is an isomorphism of functors from $\mathrm{D}(\mathrm{Qcoh}S)$ to $\mathrm{D}(\bar{F})$ [7, (3.18)]. We now have

$$\begin{aligned} 0 &= u^* \mathrm{RHom}_S(\mathcal{T}, \mathcal{M}) \cong u^* R p_* \mathrm{RHom}_S(\mathcal{T}, \mathcal{M}) \quad \text{by [7, p.85]} \\ &\cong R q_* v^* \mathrm{RHom}_S(\mathcal{T}, \mathcal{M}) \quad \text{by [7, (3.18)]} \\ &\cong R q_* v^* (\mathcal{T}^\vee \otimes_S^L \mathcal{M}) \\ &\cong R q_* (\bar{\mathcal{T}}^\vee \otimes_{\bar{S}}^L L v^* \mathcal{M}) \\ &\cong R q_* \mathrm{RHom}_{\bar{S}}(\bar{\mathcal{T}}, L v^* \mathcal{M}) \\ &\cong \mathrm{RHom}_{\bar{S}}(\bar{\mathcal{T}}, L v^* \mathcal{M}). \end{aligned}$$

But $\bar{\mathcal{T}}$ generates $\mathrm{D}(\mathrm{Qcoh}\bar{S})$ so $v^* \mathcal{M} = 0$. Since v^* is faithful, $\mathcal{M} = 0$, and we are done. \square

Corollary 4.6 (cf. [3], Corollary 5.2). *The functor $\mathrm{Hom}_S(\mathcal{T}, -) : \mathrm{coh}(S) \rightarrow \mathrm{mod}A$ induces an isomorphism*

$$\mathrm{Hom}_S(\mathcal{T}, -) : K_*(S) \rightarrow K_*(F \times B \times Q).$$

Proof. It follows from Theorem 1.98 of [10] that the equivalence of derived categories found in Theorem 4.5 induces an isomorphism in K -theory

$$\mathrm{Hom}_S(\mathcal{T}, -) : K_*(\mathrm{coh}S) \rightarrow K_*(\mathrm{mod}A).$$

Moreover, A has a nilpotent ideal I so that A/I is isomorphic to its semi-simple quotient $F \times B \times Q$. Thus, it follows that the K -theory of A is isomorphic to that of $F \times B \times Q$, and we recover the isomorphism found in [3]. \square

REFERENCES

- [1] M. Auslander, I. Reiten, and S. Smalø, *Representation Theory of Artin Algebras*, Camb. Studies in Adv. Math., No. 36, 1995.
- [2] D. Baer, Tilting sheaves in representation theory of algebras, *Manus. Math.*, **60** (1988) 323-347.
- [3] M. Blunk, Del Pezzo surfaces of degree 6 over an arbitrary field, *J. Algebra*, **323** (2010), no. 1, 42–58.
- [4] A. Bondal and M. Van den Bergh, Generators and representability of functors in commutative and noncommutative geometry, *Moscow Mathematical Journal*, **3** (2003), no. 1, 1-36.
- [5] D.A. Cox, The homogeneous coordinate ring of a toric variety, *J. Alg. Geom.*, **4** (1995), no. 1, 17-50.
- [6] L. Hille and M. Van den Bergh, Fourier-Mukai transforms, pp. 147-173 in *Handbook of Tilting Theory*, Lond. Math. Soc. Lect. Note Series (No. 332), Eds. L.A. Hügel, D. Happel, and H. Krause, Camb. Univ. Press, 2007.
- [7] D. Huybrechts, *Fourier-Mukai transforms in algebraic geometry*, Oxford Univ. Press, 2006.
- [8] A. Neeman, The connection between the K-theory localization theorem of Thomason, Trobaugh and Yao and the smashing subcategories of Bousfield and Ravenel, *Ann. Sci. de l'Éc. Norm. Sup.*, **25** (1992), no. 5, 547-566.
- [9] S.P. Smith, Computation of the Grothendieck and Picard groups of a toric DM stack \mathcal{X} by using a homogeneous coordinate ring for \mathcal{X} , *Glasgow Math. Jour.*, to appear. arXiv:0806.0192v2.
- [10] R. W. Thomason and T. Trobaugh, Higher algebraic K-theory of schemes and of derived categories, pp. 247–435 in *The Grothendieck Festschrift, Vol. III*, Progr. Math., **88**, Birkhäuser Boston, Boston, MA, 1990.

DEPARTMENT OF MATHEMATICS, UNIV. OF BRITISH COLUMBIA., DEPARTMENT OF MATHEMATICS,
BOX 354350, UNIV. WASHINGTON, SEATTLE, WA 98195

E-mail address: mblunk@math.ubc.ca, sjsierra@math.washington.edu, smith@math.washington.edu